

# Virginia Medicaid Managed Care

## Behavioral Health HEDIS 2020

Performance Measure	Virginia State Rate	National 50 <sup>th</sup> Percentile Rate
<b>Follow-Up Care for Children Prescribed ADHD Medication</b>		
The percentage of children 6-12 years of age newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.		
Initiation Phase	38.05%	43.05%
Continuation & Maintenance Phase	NA*	54.77%
<b>Antidepressant Medication Management</b>		
The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment.		
Effective Acute Phase Treatment	57.40%	53.65%
Effective Continuation Phase Treatment	44.21%	38.37%
<b>Follow-Up After Hospitalization for Mental Illness</b>		
The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider.		
7 day follow-up total	33.39%	36.78%
30 day follow-up total	56.62%	59.38%
<b>Follow-Up After Emergency Department Visit for Mental Illness</b>		
The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness.		
7 day follow-up total	44.89%	38.98%
30 day follow-up total	59.97%	55.15%
<b>Adherence to Antipsychotic Medications for Individuals with Schizophrenia</b>		
The percentage of members 18 years of age and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.		
	62.66%	62.06%
<b>Diabetes Screening for People with Schizophrenia or Bipolar who are Using Antipsychotic Medications</b>		
The percentage of members 18–64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.		
	82.26%	82.09%

\*Note: Virginia measure rates performing at or above the 50<sup>th</sup> percentile are represented in green.

\*Definition:

Not Applicable (NA): is displayed when the sample size is not large enough or if the HEDIS rate measures are not reported using a 95 percent confidence interval.

# Medallion 4.0

## Behavioral Health HEDIS 2020

### Performance by Managed Care Organization

Performance Measure	Aetna	Anthem	Magellan	Optima	United	Virginia Premier	National 50 <sup>th</sup> Percentile Rate
<b>Follow-Up Care for Children Prescribed ADHD Medication</b>							
The percentage of children 6-12 years of age newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.							
Initiation Phase	53.61%	42.49%	30.00%	37.96%	NA*	28.04%	43.05%
Continuation & Maintenance Phase	76.92%	59.59%	NA*	52.36%	NA*	49.15%	54.77%
<b>Antidepressant Medication Management</b>							
The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment.							
Effective Acute Phase Treatment	55.60%	51.39%	58.36%	55.20%	52.69%	58.29%	53.65%
Effective Continuation Phase Treatment	40.39%	34.92%	50.93%	42.00%	42.30%	41.77%	38.37%
<b>Follow-Up After Hospitalization for Mental Illness</b>							
The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider.							
7 day follow-up total	35.42%	43.41%	23.69%	40.49%	36.03%	40.07%	36.78%
30 day follow-up total	58.97%	64.27%	43.46%	64.70%	58.01%	61.68%	59.38%
<b>Follow-Up After Emergency Department Visit for Mental Illness</b>							
The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness.							
7 day follow-up total	42.40%	53.11%	41.05%	53.39%	41.50%	48.85%	38.98%
30 day follow-up total	57.60%	65.03%	56.17%	63.56%	54.75%	61.55%	55.15%
<b>Adherence to Antipsychotic Medications for Individuals with Schizophrenia</b>							
The percentage of members 18 years of age and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.							
	51.42%	52.33%	59.32%	49.08%	39.13%	46.92%	62.06%
<b>Diabetes Screening for People with Schizophrenia or Bipolar who are Using Antipsychotic Medications</b>							
The percentage of members 18–64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.							
	83.68%	78.31%	81.58%	72.28%	82.86%	83.07%	82.09%

\* Definition: Not Applicable (NA): is displayed when the sample size is not large enough or if the HEDIS rate measures are not reported using a 95 percent confidence interval.

# Commonwealth Coordinated Care (CCC Plus)

## Behavioral Health HEDIS 2020

### Performance by Managed Care Organization

Performance Measure	Aetna	Anthem	Magellan	Optima	United	Virginia Premier	National 50 <sup>th</sup> Percentile Rate
<b>Follow-Up Care for Children Prescribed ADHD Medication</b>							
The percentage of children 6-12 years of age newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.							
Initiation Phase	56.57%	46.19%	14.77%	37.20%	37.33%	10.50%	43.05%
Continuation & Maintenance Phase	NA*	53.45%	NA*	53.57%	NA*	16.67%	54.77%
<b>Antidepressant Medication Management</b>							
The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment.							
Effective Acute Phase Treatment	51.47%	51.82%	60.67%	53.55%	67.39%	67.52%	53.65%
Effective Continuation Phase Treatment	38.99%	39.19%	47.78%	40.74%	53.86%	54.66%	38.37%
<b>Follow-Up After Hospitalization for Mental Illness</b>							
The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider.							
7 day follow-up total	30.56%	32.25%	18.69%	34.09%	31.03%	32.31%	36.78%
30 day follow-up total	56.67%	56.96%	39.89%	60.54%	55.83%	56.52%	59.38%
<b>Follow-Up After Emergency Department Visit for Mental Illness</b>							
The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness.							
7 day follow-up total	42.76%	43.97%	47.48%	45.49%	39.61%	44.47%	38.98%
30 day follow-up total	62.76%	62.76%	63.21%	62.01%	57.46%	61.65%	55.15%
<b>Adherence to Antipsychotic Medications for Individuals with Schizophrenia</b>							
The percentage of members 18 years of age and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.							
	69.61%	68.21%	71.05%	66.76%	63.43%	69.76%	62.06%
<b>Diabetes Screening for People with Schizophrenia or Bipolar who are Using Antipsychotic Medications</b>							
The percentage of members 18–64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.							
	82.12%	84.50%	80.57%	78.12%	84.78%	87.13%	82.09%

\*Definition: Not Applicable (NA): is displayed when the sample size is not large enough or if the HEDIS rate measures are not reported using a 95 percent confidence interval.